
This Directed Acyclic Graph and write-up is an excerpt from a larger NASA document.

## NASA/TP-20220015709

Directed Acyclic Graphs: A Tool for Understanding the NASA Spaceflight Human System Risks

**Human System Risk Board** 

October 2022



## Microhost Risk DAG Narrative

- The Microhost Risk centers around the possibility for microbial contamination leading to Infections that if left inadequately treated could become Sepsis. Both Infections and Sepsis can lead to deterioration of Individual Readiness and Crew Capability which affects Task Performance, likelihood of Evacuation for medical reasons, and in severe cases can contribute to Loss of Crew Life. They can also lead to Long Term Health Outcomes if inadequately treated and post-mission/career Surveillance enables Detection of Long Term Health Outcomes to understand the magnitude of the problem.
- The cause of infections can come from various sources:
  - Microbial Virulence Factors evidence that the virulence of certain microbes change in response to spaceflight environment.
  - This may lead to an increased risk of infections.
  - Can indirectly lead to infections through changes in Microbiome.
  - Immune (Risk) the strength of the immune system determines how well individuals fight
    off infections.
  - Surface Contamination microbes on surfaces are found regularly on ISS, cleaning procedures can decrease impact on crew.
  - Air Contamination good air quality and filtration can limit likelihood of airborne and dropletbased infections among crew.
  - Water Contamination water quality monitoring and cleaning helps limit infections in crew.
  - Pharmaceutical Contamination repackaged pharmaceuticals are susceptible to contamination increasing risk for infection among crew.
  - Food Contamination inadequate packaging and storage conditions for crew food could lead to infections including gastroenteritis.
- Countermeasures that affect microbial levels must be included in the Crew Health and
  Performance System and accommodated in Vehicle Design. These are affected by the HSIA
  (Risk) and include: Countermeasures include the storage conditions which if compromised could
  increase contamination of food and pharmaceuticals; the storage conditions are also impacted by
  the food system available which is represented in the DAG by the Food and Nutrition (Risk).
  - Preventive Source Control includes monitoring, regular cleaning, filtration and other modes
    of limiting spread of microbes.
  - Hygiene includes personal hygiene such as regular showers, dental hygiene, and other personal cleaning that limits the development of Infection.
  - Environmental Monitoring Capability is necessary to Detect Contamination levels in the air, water, and surfaces. This enables Intervention Source Control measures like cleaning or maintenance of filtration systems.
  - Medical Treatment Capability includes antibiotics, antifungal, and antiviral medications, as well as other supportive care, intended to minimize consequence of infection and prevent the development of sepsis.

-	<b>Infections</b> and <b>Sepsis</b> affect cognitive function, mood and performance and therefore affect <b>Behavioral (Risk)</b> and <b>Team (Risk)</b> which negatively impacts <b>Individual Readiness</b> and <b>Crew Capability.</b>