

This document begins with four LRL photos of sample 12034 and concludes with the 12034 Lunar Sample Compendium page.

Sample 12034 is a regolith breccia, formed from soil compacted in an impact. Pete Conrad ran into the rock while he was digging a trench 15 meters inside the rim of Head Crater. Interestingly, characteristics of the compacted soil do not match those of the soil taken from the trench, indicating that sample 12034 was thrown onto the site from a non-mare site and then was buried under about 15 cm of mare soil thrown onto the area from a third, mare site.



LRL photo S69-60934



LRL photo 60939



LRL photo S69-60943



LRL photo S69-60945

12034
Regolith Breccia
155 grams

DRAFT




Figure 1: Lunar sample 12034 after first saw cut in 1970. NASA S75-34235. Cube is 1 cm.

Introduction

12034 was collected from the bottom of the same trench as soil sample 12033. These samples were located about 15 meters inside the rim of Head Crater (Schoemaker et al. 1970). *“Let’s sample that rock that I dug up from down deep (~15 cm). Let me get a photo of it (AS12-49-7195 - 6)”*.


12034 is a regolith breccia, but it is largely made up of components foreign to the Apollo 12 regolith and is from some distant, non-mare, regolith (figure 1). It

contains abundant KREEP; both as glasses and annealed lithic fragments.

12034 has not been dated, but there is evidence in the U/Pb data that there has been a young event.

Petrography

Simon et al. (1985) described regolith breccias from Apollo 12 and give new analyses. They note that although 12034 is a soil breccia, it is unlike the soils at



Mode for 12034

	Chao et al. 1971	McGee et al. 1979
Basaltic rock	5.2 %	5.96
Anorthositic rock	1.6	
Mineral fragments	14.7	12
Glass-welded aggregate	25.7	
Devitrified glass	2.2	
Heterogeneous glass	6.6	23
Homogeneous glass	0.3	tr.
Basaltic microbreccia	0.3	5
Anorthositic microbx.	1.8	
Less than 25 microns	27.3	53
Pore space	14.2	

Figure 2: Photo of thin section 12034,106 showing clastic texture. Scale is about 1.2 cm.




Figure 3: Thin section photomicrograph of 12034. NASA S70-49463. Scale unknown.

Modal Petrology for 12034

	Simon et al. 1985
Mare Basalt	1.7
ANT	3
CMB	0.7
Poik. Bas.	1.2
Regolith bx.	2.8
Agglutinate	8
Pyroxene	1.5
Olivine	0.3
Plag.	2
Opagues	
Glass	5.2
Matrix	45.5

Apollo 12. Simon et al. (1985) and Anderson and Smith (1971) note that it contains Mg-rich orthopyroxene and olivine.

Sample 12034 is a fragmental matrix breccia similar in appearance to the Apollo 11 fragmental matrix breccias, but with a much lower porosity (figure 2). According to McGee et al. (1979) it consists of a variety of glass, mineral and lithic clasts contained in a matrix of brown glass fragments and comminuted debris (figure 3). The texture of the matrix is seriate with fragments ranging in size from the limit of resolution up to 2 mm. Brown glass fragments dominate the less than 0.05 mm size fraction (Phinney et al. 1976). Waters et al. (1971) noted a vague stratified nature of the sample. Chao et al. (1971) and Anderson and Smith (1971) reported 14% and 10% pore space, respectively.

The glass clast population in 12034 is extremely diverse, ranging from colorless, pale green, yellow or orange homogeneous glass to completely devitrified glass with included mineral grains (Chao et al. 1971; von Engelhardt et al. 1971). Subrounded to rounded fragments of maskelynite and devitrified maskelynite occur commonly as inclusions in the large devitrified glass clasts. Devitrification features in glass clasts take the form of vaiolitic clusters of plagioclase needles and more commonly axiolitic intergrowths of tightly packed plagioclase and pyroxene crystals (McGee et al. 1979).

The lithic clasts in 12034 include basalts, dark-matrix breccia, cataclastic anorthosite and anorthositic gabbro (see modes in tables). Rare basalts have textures ranging from ophitic to vitrophyric. Anderson and Smith (1971) reported “grey mottled” basalts in 12034. These are made up of about half plagioclase and half low-Ca pyroxene (orthopyroxene?) and have the bulk




Figure 4: Composition of pyroxene found in matrix of 12034.




Figure 5: Comparison of REE for three regolith breccias from Apollo 12 (data from Goles et al. 1971 and Wanke et al. 1971).




Figure 6: Composition of regolith breccia compared with all lunar soils (see table).

composition of KREEP. Based on what we know now, they are probably fragments of impact melt.

Chemistry

One can use the whole rock U and Th measurements from radiation counting (O’Kelley et al. 1971,

Rancitelli et al. 1971) to judge which analyses are representative of the sample (table 1). The split obtained by Simon et al. (1985) is high in U and Th, while the splits analyzed by Goles et al. (1971) and Wakita et al. (1971) appear low.

Using chemical mixing models, but with different assumed compositions for end members, Meyer et al. (1971) calculated that 12034 is ~30 % mare basalt, ~5 % anorthosite and ~ 66 % KREEP, while Simon et al. (1985) calculated that 12034 is ~ 1 % basalt, 8 % anorthosite and 89 % low-KREEP. Finally, McKay et al. (1971) calculated ~ 56 % KREEP for 12034, using K, Rb and Ba only.

Radiogenic age dating

Tatsumoto et al. (1971) determined the U, Th and Pb isotopes in 12034, providing evidence for a young event (figure 7). The Nd – Sm data give a model age for KREEP in 12034 of 4.3 – 4.4 b.y. (Lugmair and Carlson 1978).

Cosmogenic isotopes and exposure ages

O’Kelly et al. (1971) determined the cosmic ray induced activity of 12034 as $^{22}\text{Na} = 29$ dpm/kg, $^{26}\text{Al} = 45$ dpm/kg, $^{46}\text{Sc} = <10$ dpm/kg, $^{48}\text{V} = <60$ dpm/kg, $^{54}\text{Mn} = 16$ dpm/kg, $^{56}\text{Co} = <16$ dpm/kg and $^{60}\text{Co} = <4$ dpm/kg. Rancitelli et al. (1971) determined $^{22}\text{Na} = 34$ dpm/kg, $^{26}\text{Al} = 60$ dpm/kg, $^{54}\text{Mn} = <40$ dpm/kg, $^{56}\text{Co} = <56$ dpm/kg and $^{60}\text{Co} = 1.5$ dpm/kg. Breccia 12034 was shielded from solar cosmic rays, having been buried in a trench.

Basford et al. (1973) determined the isotopic composition of Kr and Xe, but do not give an exposure age.

Other Studies

- Epstein and Taylor (1971) oxygen isotopes
- Reed and Jovanovic (1971) halogens

Processing

One end was cut off 12034 in 1970, and in 1975, a small slab was cut (see figures). There are 16 thin sections. 12034 was never studied in “consortium mode”.




Figure 7: U/Pb concordia diagram showing that 12034 data is discordant (Tatsumoto et al. 1971).




Table 1. Chemical composition of 12034.

reference weight	Simon85	Wakita 71 0.199 .267 g		O'Kelly71 155 g	Hubbard73	Wiesmann75	Goles71	Rancitelli71	Tatsumoto71
SiO2 %		45.8		(a)	47.76 (d)		45.78 (e)		
TiO2	1.8 (e)	2.7	2.8	(a)	2.33 (d)	2.28 (c)	2.3 (e)		
Al2O3	14.7 (e)	15.7	15.7	(a)	15.49 (d)		14.6 (e)		
FeO	12 (e)	13.3		(a)	12.42 (d)		13.29 (e)		
MnO	0.156 (e)	0.167	0.173	(a)	0.19 (d)		0.17 (e)		
MgO	8.3 (e)	9.6		(a)	8.31 (d)				
CaO	11.5 (e)	11.2	11.6	(a)	10.85 (d)		11.3 (e)		
Na2O	0.78 (e)	0.788	0.755	(a)	0.67 (d)	0.86 (c)	0.78 (e)		
K2O	0.67 (e)		0.529	(a) 0.55 (b)	0.48 (d)	0.5 (c)		0.56 (b)	
P2O5					0.53 (d)				
S %					0.09 (d)				
sum									
Sc ppm	29 (e)	31		(a)			30 (e)		
V	65 (e)	90	110	(a)					
Cr	1724 (e)	35		(a)		1790 (c)	1840 (e)		
Co	17.5 (e)						30.4 (e)		
Ni	90 (e)								
Cu									
Zn									
Ga									
Ge ppb									
As									
Se									
Rb			11 (a)			12.75 (c)			
Sr	170 (e)					195 (c)			
Y			186 (a)						
Zr	870 (e)	640		(a)		756 (c)	630 (e)		
Nb									
Mo									
Ru									
Rh									
Pd ppb									
Ag ppb									
Cd ppb									
In ppb			35 (a)						
Sn ppb									
Sb ppb									
Te ppb									
Cs ppm			0.48 (a)						
Ba	850 (e)	470		(a)		748 (c)	720 (e)		
La	68.3 (e)	75	60 (a)			68.3 (c)	60.4 (e)		
Ce	170 (e)		172 (a)			172 (c)	176.7 (e)		
Pr			20.4 (a)						
Nd	110 (e)		100 (a)			105 (c)	92 (e)		
Sm	29.5 (e)	32	29.4 (a)			29.5 (c)	28.3 (e)		
Eu	2.8 (e)	2.95	2.65 (a)			2.81 (c)	2.69 (e)		
Gd			33.3 (a)			35.6 (c)			
Tb	6.6 (e)		5.51 (a)				8.12 (e)		
Dy	42 (e)		34 (a)			38 (c)			
Ho			8.8 (a)						
Er	3.7 (e)		21.7 (a)			23.4 (c)			
Tm			3.6 (a)						
Yb	25 (e)	25	22.4 (a)			21.5 (c)	21.7 (e)		
Lu	3.6 (e)	3.3	2.74 (a)				3.14 (e)		
Hf	22 (e)	27		(a)		20.4 (c)	20.4 (e)		
Ta	3.3 (e)						3.09 (e)		
W ppb									
Re ppb									
Os ppb									
Ir ppb									
Pt ppb									
Au ppb									
Th ppm	17.2 (e)	13.2		(a) 13.1 (b)		11.7 (c)	11.53 (e)	13.9 (b)	13 (c)
U ppm	4.5 (e)			3.4 (b)		3.4 (c)		3.53 (b)	3.5 (c)

technique (a) RNAA, (b) radiation counting, (c) IDMS, (d) XRF, (e) INAA

76671

DRWG COMPLETED
JUNE 2, 70







Figure 8: Group photo of 12034 after cutting slab (.82) and end (.80-.84) pieces in 1975.
 NASA S75-34414. Cube is 1 cm.






Figure 9: Sawn surface of middle piece (,9) of 12034 - side facing slab (,82). Cube is 1 cm. NASA S75-34243.




Figure 10: Processing photo of 12034 after cutting end ,80. Location of slab ,82 is approximate.



Figure 11: Slab (82) cut from 12034 facing middle piece (9). Cube is 1 cm. NASA S75-34238.

References for 12034

- Anderson A.T. and Smith J.V. (1971) Nature occurrence and exotic origin of "grey mottled" (Luny Rock) basalts in Apollo 12 soils and breccias. Proc. 2nd Lunar Sci. Conf. 431-438.
- Basford J.R., Dragon J.C., Pepin R.O., Coscio M.R. and Murthy V.R. (1973) Krypton and Xenon in lunar fines. Proc. 4th Lunar Sci. Conf. 1915-1955.
- Chao E.C.T., Boreman J.A. and Desborough G.A. (1971) The petrology of unshocked and shocked Apollo 11 and Apollo 12 microbreccias. Proc. 2nd Lunar Sci. Conf. 797-816.
- von Engelhardt W., Arndt J., Muller W.F. and Stoffler D. (1971) Shock metamorphism and origin of regolith and breccias at the Apollo 11 and Apollo 12 landing sites. Proc. 2nd Lunar Sci. Conf. 833-854.
- Epstein S. and Taylor H.P. (1971) O18/O16, Si30/Si28, D/H and C13/C12 ratios in lunar samples. Proc. 2nd Lunar Sci. Conf. 1421-1441.
- Ferland R.M. (1983) Regolith Breccia Workbook. JSC 19045
- Goles G.G., Ducan A.R., Lindstrom D.J., Martin M.R., Beyer R.L., Beyer R.L., Osawa M., Randle K., Meek L.T., Steinborn T.L. and McKay S.M. (1971) Analysis of Apollo 12 specimens: Compositional variations, differentiation processes and lunar soil mixing models. Proc. 2nd Lunar Sci. Conf. 1063-1081.
- LSPET (1970) Preliminary Examination of Lunar Samples from Apollo 12 Science 167, 1325-1339
- Lugmair G.W. and Carlson R.W. (1978) The Sm-Nd history of KREEP. Proc. 9th Lunar Planet. Sci. Conf. 689-704.




Figure 12: Slab (,82) cut from 12034 side facing (,80) end. NASA S 75-34236. Cube is 1 cm.

McGee P.E., Simonds C.H., Warner J.L. and Phinney W.C. (1979) Introduction to the Apollo Collections: Part II Lunar Breccias. JSC Curator's Office

McKay D.S., Morrison D.A., Clayton U.S., Ladle G.H. and Lindsay J.F. (1971) Apollo 12 soils and breccias. Proc. 2nd Lunar Sci. Conf. 755-773.

Meyer C., Brett R., Hubbard N.J., Morrison D.A., McKay D.S., Aitken F.K., Taketda H. and Schonfeld E. (1971) Mineralogy, chemistry and origin of the KREEP component in soil samples from the Ocean of Storms. Proc. 2nd Lunar Sci. Conf. 393-411.

O'Kelly, G.D., Eldridge J.S., Schonfeld E. and Bell P.R. (1971) Abundances of the primordial radionuclides K, Th and U in Apollo 12 samples. Proc. 2nd Lunar Sci. Conf. 1159-1168.

O'Kelly, G.D., Eldridge J.S., Schonfeld E. and Bell P.R. (1971b) Cosmogenic radionuclide concentrations and

exposure ages of lunar samples from Apollo 12 samples. Proc. 2nd Lunar Sci. Conf. 1747-1755.

Phinney W.C., McKay D.S., Simonds C.H. and Warner J.L. (1976) Lithification of vitric- and clastic-matrix breccias: SEM petrography. Proc. 7th Lunar Sci. Conf. 2469-2493.

Rancitelli L.A., Perkins R.W., Felix W.D. and Wogman N.A. (1971) Erosion and mixing of the lunar surface from cosmogenic and primordial radionuclide measurements in Apollo 12 samples. Proc. 2nd Lunar Sci. Conf. 1757-1772.

Reed G.W. and Jovanovic S. (1971) The halogen and other trace elements in Apollo 12 samples. Proc. 2nd Lunar Sci. Conf. 1261-1276.

Schoemaker E.M., Batson R.M., Bean A.L., Conrad C., Dahlem D.H., Goddard E.N., Hait M.H., Larson K.B., Schaber G.C., Schleicher D.L., Sutton R.L., Swann G.A., and Waters A.C. (1970) 10. Preliminary Geologic Investigation of the Apollo 12 landing site. In Apollo 12: Preliminary Science Rpt. NASA SP-215. 113-156.




Figure 13: Sawn surface of 12034,80 (butt end). NASA S75-34244.

Simon S.B., Papike J.J. and Gosselin D.C. (1985) Petrology and chemistry of Apollo 12 regolith breccias. Proc. 16th Lunar Planet. Sci. Conf. D75-86.

Tatsumoto M., Knight R.J. and Doe B.R. (1971) U-Th-Pb systematics of Apollo 12 lunar samples. Proc. 2nd Lunar Sci. Conf. 1521-1546.

Wakita H. and Schmitt R.A. (1971) Bulk elemental composition of Apollo 12 samples: Five igneous and one breccia rocks and four soils. Proc. 2nd Lunar Sci. Conf. 1231-1236.

Wakita H., Rey P. and Schmitt R.A. (1971) Abundances of the 14 rare-earth elements and 12 other trace elements in Apollo 12 samples. Proc. 2nd Lunar Sci. Conf. 1319-1329.

Waters A.C., Fisher R.V., Garrison R.E. and Wax D. (1971) Matrix characteristics and origin of lunar breccia samples 12034 and 12073. Proc. 2nd Lunar Sci. Conf. 893-907.

Wiesmann H. and Hubbard N.J. (1975) A compilation of the Lunar Sample data generated by the Gast, Nyquist and Hubbard PI-ships. JSC unpublished